

# NH<sub>3</sub> emissions from grazing pasture following urea and urease inhibitor treatments

Mei Bai<sup>1\*</sup>, Helen Suter<sup>1</sup>, Shu Kee Lam<sup>1</sup>, Rohan Davies<sup>2</sup>, Deli Chen<sup>1</sup>

<sup>1</sup> Crop and Soil Sciences Section, Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, VIC 3010, Australia

<sup>2</sup>BASF Australia Ltd., Southbank, VIC 3006, Australia

\*Corresponding author: [mei.bai@unimelb.edu.au](mailto:mei.bai@unimelb.edu.au)

## Abstract

Ammonia (NH<sub>3</sub>) volatilization to the atmosphere following urea based nitrogen (N) fertilizer application not only causes nutrient loss but also detrimentally impacts on the environment, our ecosystems, and contributes to global warming (as an indirect greenhouse gas). Here, we report our studies of quantifying NH<sub>3</sub> emissions from dairy pasture following urea application, and the effectiveness of a urease inhibitor in mitigating NH<sub>3</sub> loss. Two experiments were conducted in summer at Queensland (northern site) and autumn at South Australia (southern site) where urea was surface applied to pasture. A urease inhibitor (NBPT, applied with urea as Green ureaNV<sup>TM</sup>) was added at the northern site. Open-path NH<sub>3</sub> laser concentration sensors were used to measure line-averaged concentrations along an open path downwind of the treatment plots. Ammonia fluxes were calculated using the inverse-dispersion technique (WindTrax). We found NH<sub>3</sub> flux increased following urea application and varied temporally at the two sites. Daily average NH<sub>3</sub> flux from dairy pastures fertilized with urea was  $4.4 \pm 0.46$  and  $6.4 \pm 1.2$  mg N m<sup>-2</sup> h<sup>-1</sup> for the northern and southern sites, respectively. Nitrogen loss as volatilised NH<sub>3</sub> from the urea application over the course of the experiments (12-15 days) accounted for approximately 40 and 60% of total applied N for the northern and southern sites, respectively. The difference between sites is likely attributed to the differences in N input, soil properties and microbial activity. The urease inhibitor reduced NH<sub>3</sub> emissions by approximately 71% compared to that from the urea treatment. The results in these studies also demonstrated that inverse-dispersion technique combined with the open-path lasers is able to measure NH<sub>3</sub> fluxes from large-scale field sites, and the open-path NH<sub>3</sub> laser has adequate detection resolution.

**Key Words** Open-path lasers, NH<sub>3</sub> emissions, grazing pasture, micrometeorological inverse-dispersion technique, urea fertilizer

## Introduction

Applying nitrogen (N) fertilizer to grazing pasture is a common practice on dairy farms. Urea fertilizer applied to the soil surface can be hydrolysed rapidly to ammonia (NH<sub>3</sub>). The NH<sub>3</sub> volatilization to the atmosphere can be a significant loss of valuable N from the fertilizer, and has negative impacts on the environment and human health, through the formation of aerosols (Sutton et al., 1998). In addition, the depositions of NH<sub>3</sub> downwind of farm can result in the emission of nitrous oxide (N<sub>2</sub>O), a potent greenhouse gas, through the process of nitrification and denitrification, and eutrophication of waterways, consequently decreasing biodiversity in terrestrial ecosystems (Mosier et al., 1998; Smith et al., 2007). Application of fertiliser amendments has been used to reduce NH<sub>3</sub> emissions, e.g., urease inhibitor along with urea fertilizer is one of the options for increasing N fertilizer efficiency. It is well known that the urease inhibitor can control the hydrolysis of urea by inhibiting the urease enzyme activity (Chen et al., 2008).

Numerous studies on quantifying NH<sub>3</sub> emissions after N-fertilizer application have been conducted. Chamber techniques (e.g. wind tunnel) have been used to measure NH<sub>3</sub> emissions (Lockyer, 1984). The principle of chamber techniques is to draw air from a tunnel (cover above the fertilizer treated soil) to a flask which is filled with acidic solution that traps the NH<sub>3</sub>, and then the NH<sub>3</sub> flux is determined by the calculation of NH<sub>3</sub> absorption. Micrometeorological techniques have the advantages of being a non-intrusive approach and representing real environmental conditions, e.g. the integrated horizontal flux (Denmead et al., 1998), and the passive sampling techniques (Leuning et al., 1985). The latter techniques are widely used in field measurements. However, the passive sampling techniques are labour intensive and have a low resolution of continuous measurements. Laubach et al. (2012) reported that the backward turbulence contributed to the uncertainty of flux measurement using this sampling technique. In the last two decades, a model based inverse-dispersion technique has become well-established with the simplicity of field flux measurements from a well-defined source area given by a single height concentration and wind statistics (Flesch et al.,

1995). Furthermore, line-average  $\text{NH}_3$  concentration sensors (e.g., lasers) have been commonly used in the large-scale measurements (up to 500 m path length), and no pump and tubing are required.

Here we report on two field studies to measure  $\text{NH}_3$  emissions from dairy pasture following urea application at two locations (northern and southern sites) in summer and autumn using inverse-dispersion technique combined with open-path  $\text{NH}_3$  concentration sensors. Furthermore, we added a urease inhibitor in conjunction with urea application at the northern site to investigate  $\text{NH}_3$  reduction compared to  $\text{NH}_3$  emissions with urea treatment.

## Methods

### *Experimental site*

#### *Northern site*

The first experiment was conducted at a research farm (27.54°S, 152.34°E), at the University of Queensland Gatton campus, 80 km west of Brisbane, Queensland from 23 November to 12 December 2013. The terrain was open and flat with short grass. The soil is typical silt loam, with a pH of 5.9 and 2.4% organic carbon (Table 1). No animals were grazed in this area. Two experimental plots (15 m radius) were constructed for urea treatment (U) and urea + urease inhibitor (NBPT) treatment (GU). The distance between these two plots was 50 m. Urea was applied to U plot surface at rate of 40 kg N ha<sup>-1</sup> on 26 November 2013 and Green ureaNV™ (40 kg N ha<sup>-1</sup>, containing urea and urease inhibitor) was applied to GU plot surface on the same day. Ammonia measurements were made prior to the application and ended on 12 December. During the measurement period, the average maximum and minimum temperature was 30.2 and 16.0°C, respectively. The total rainfall was 79.4 mm (BOM, 2016a).

Two open-path  $\text{NH}_3$  concentration sensors (NEO Monitors LaserGas, Norsk Elektro Optikk, Norway) were deployed using tuneable diode laser absorption spectroscopy technique. A transmitter combined with a single corner cube retro reflector measures the line-average  $\text{NH}_3$  concentrations along the open path. The detection limit for  $\text{NH}_3$  was less than 10 ppb at 100 m one way with response time of 1–2 seconds. One concentration sensor was placed 10 m west of U and GU plots separately, with each path length 80 m between transmitter and retro reflector at a height of 1.5 m above ground. One minute averaged concentrations of  $\text{NH}_3$  were recorded. A weather station was set up 100 m south of the plots coupled with a three-dimensional (3-D) sonic anemometer (CSAT3, Campbell Scientific, Logan, UT, USA) at a height of 2.0 m above ground. Ten-minute average wind speed, wind direction, air temperature and air pressure were collected by a data logger (CR23X, Campbell Scientific, Logan, UT, USA) at a frequency of 10 Hz.

#### *Southern site*

The second experiment was conducted at a commercial dairy farm (38°01'S 140°55'E), 27 km southeast of Mount Gambier, South Australia between September and October 2014. The topography of the experimental site was open and flat and was covered by short grass. The soil is classified as clay loam, with a pH of 4.9 and 4.9% organic carbon (Table 1). There were no animals on site. One 25 m-radius experimental circle was created. Urea was spread to the pasture surface at rate of 50 kg N ha<sup>-1</sup> on 30 September 2014. Measurements were made prior to the application and ended on 23 October. During the experimental period, the average maximum and minimum temperature were recorded at 18.6 and 8.0°C, respectively. The total rainfall was 63.4 mm and average wind speed at 3 pm was 4.9 m s<sup>-1</sup> (BOM, 2016b). The predominant winds during the study period were south-west-southwest.

Two open-path  $\text{NH}_3$  Neo laser concentration sensors were used. One sensor was placed on the southern side and the other one was placed on the northern side of the plot; both were 5 m to the edge of the plot at a height of 1.48 m above ground. One minute average  $\text{NH}_3$  concentrations were measured with the path length of 116 and 78 m for the laser on the north side and south side, respectively. A weather station coupled with a 3-D sonic anemometer (CSAT3, Campbell Scientific, Logan, UT, USA) was set up at a height of 2.47 m above ground, 300 m on the west side of the plot.

### *Data processing*

At both sites two  $\text{NH}_3$  concentration sensors were run side-by-side to calibrate the difference between the two sensors for two days prior to the flux measurements. Ammonia concentrations were not used for the flux calculation when light intensity of the concentration sensor was less than 10%. The wind turbulent parameters including wind friction velocity  $u_*$  (m s<sup>-1</sup>), Obukhov stability length  $L$  (m), surface roughness length  $z_0$  (m) and wind statistics were derived from sonic measurements. Following the filtering procedures

(Flesch et al., 2014), the error-prone data ( $u_* < 0.05 \text{ m s}^{-1}$ ,  $|L| < 5 \text{ m}$  and  $z_0 > 0.5 \text{ m}$ ) were removed. Valid concentrations and wind variables were merged to 10-minute averages using SAS software (SAS 9.3, SAS Institute Inc., Cary, NC, USA). Ammonia fluxes from the experimental plot were calculated using the inverse-dispersion technique (WindTrax version 2.0.8.8, Thunder beach Scientific) given the line-averaged concentrations, wind turbulent statistics and defined source area.

## Results

### Northern site fluxes

Ten-minute average  $\text{NH}_3$  fluxes show a temporal variation over the course of the experiment following urea application at the northern site (Fig 1A). As expected,  $\text{NH}_3$  fluxes showed a significant correlation with wind speed and air temperature ( $r = 0.50$ ,  $P < 0.001$ ). Less than 10% of the flux measurements were observed before 6:00 or after 18:00, when the  $\text{NH}_3$  losses were most likely negligible. The daily average  $\text{NH}_3$  flux from U and GU plots were  $4.39 \pm 0.46$  (mean  $\pm$  SE,  $n = 228$ ) and  $1.25 \pm 0.15$  (mean  $\pm$  SE,  $n = 529$ )  $\text{mg N m}^{-2} \text{ h}^{-1}$ , accounting for 39.5 and 11.3% of total applied N, respectively (Table 1). In addition, we found GU reduced  $\text{NH}_3$  volatilization by approximately 71.4% compared to U treatment. The N loss of total applied N in this study were higher than those reported on pastures in southern Victoria by Suter et al. (2013), which could be attributed to the weather conditions and soil properties.

**Table 1. Nitrogen (N) input and soil properties obtained from the northern and southern sites.**

	Northern site	Southern site
<b>N input ( kg N ha<sup>-1</sup> )</b>	Urea:40 Green UreaNV <sup>TM</sup> : 40	50
<b>Soil Property</b>		
Texture	Clay loam	Silt loam
pH (CaCl <sub>2</sub> )	5.9	4.9
Organic carbon (%)	2.4	4.9
$\delta$ CEC (meq/100g)	26.0	11.5
$\text{NH}_4^+$ -N (mg/kg)	3.5	8.9
$\text{NO}_3^-$ -N (mg/kg)	2.1	28.8


$\delta$ cation exchange capacity

**Table 2. Daily average  $\text{NH}_3$  fluxes from the northern and southern sites. Total N loss as volatilised  $\text{NH}_3$  over the measurement periods (12-15 days) at the northern and southern sites are also calculated.**


		Daily average $\text{NH}_3$ ( $\text{mg N m}^{-2} \text{ h}^{-1}$ )	N loss as volatilised $\text{NH}_3$ (%)
Northern site	Urea	$4.39 \pm 0.46$	39.5
	Green UreaNV <sup>TM</sup>	$1.25 \pm 0.15$	11.3
Southern site	Urea	$6.40 \pm 1.23$	59.5

The  $\text{NH}_3$  background concentrations for the northern site were estimated by WindTrax using the point concentrations measured by passive samplers at a height of 2.9 m above ground, 50 m on the east of the GU plot during the same measurement period (Leuning et al., 1985). The samplers were replaced twice at 9:00 and 17:00 daily, the extract from samplers was analysed at laboratory to determine the concentrations over a period of measurement, such that two concentration measurements were obtained daily. In addition, the concentrations from upwind of U plot (northwest winds) measured by open-path  $\text{NH}_3$  concentration sensor next to U plot were also used as the background concentrations of the experimental site. These estimates (samplers and concentration sensor) were adequate to reflect the variation in  $\text{NH}_3$  background concentrations.

**(A) Northern site**



**(B) Southern site**



**Fig. 1 A) Daily average NH<sub>3</sub> fluxes with urea (U) and Green UreaNV™ (GU) treatments at northern site, B) with urea treatment at southern site. Air temperatures measured at both sites are also plotted.**

*Southern site fluxes*

Daily NH<sub>3</sub> fluxes from the plot at southern site showed temperature-dependent diurnal pattern with high emission at middle of day and low emissions at night-time (Fig. 1B). Average daily NH<sub>3</sub> flux was  $6.4 \pm 1.2$  mg N m<sup>-2</sup> h<sup>-1</sup> (mean  $\pm$  SE, n = 707) over 15 days measurement period (Table 2). This fits the range of 1–10 mg N m<sup>-2</sup> h<sup>-1</sup> reported by Ni et al. (2015) using inverse-dispersion combined with FTIR techniques. Nitrogen loss as NH<sub>3</sub> volatilization over the experimental period following urea application was 59.5% of the total applied N. This value was at the higher range of values reported in the previous studies, which could be likely associated with the soil properties and fertilizer types (Zaman et al., 2008).

Greater N loss was observed at the southern site likely due to higher N input, higher  $\text{NH}_4^+$  and organic carbon contents of soils, and climatic conditions, and likely microbial activity associated with pasture species covered over the experimental site.

## Conclusions

Ammonia fluxes increased following urea application, and a urease inhibitor reduced  $\text{NH}_3$  volatilization. The inverse-dispersion technique combined with open-path  $\text{NH}_3$  concentration sensors produced reliable measurements for large-scale field studies.

## References

- BOM, 2016a. Bureau of Meteorology, Gatton, Queensland (station ID 040082) (Accessed on 14 April 2016).
- BOM, 2016b. Bureau of Meteorology, Mount Gambier, South Australia (station ID 026021) (Accessed on 14 April 2016).
- Chen, D., Suter, H., Islam, A., Edis, R., Freney, J.R., Walker, C.N., 2008. Prospects of improving efficiency of fertiliser nitrogen in Australian agriculture: a review of enhanced efficiency fertilisers. *Aust. J. Soil Res.* 46, 289-301.
- Denmead, O.T., Harper, L.A., Freney, J.R., Griffith, D.W.T., Leuning, R., Sharpe, R.R., 1998. A mass balance method for non-intrusive measurements of surface-air trace gas exchange. *Atmos. Environ.* 32, 3679-3688.
- Flesch, T.K., McGinn, S.M., Chen, D.L., Wilson, J.D., Desjardins, R.L., 2014. Data filtering for inverse dispersion emission calculations. *Agr. Forest Meteorol.* 198-199, 1-6.
- Flesch, T.K., Wilson, J.D., Yee, E., 1995. Backward-time Lagrangian stochastic dispersion models and their application to estimate gaseous emissions. *J. Appl. Meteorol.* 34, 1320-1332.
- Laubach, J., Taghizadeh-Toosi, A., Sherlock, R.R., Kelliher, F.M., 2012. Measuring and modelling ammonia emissions from a regular pattern of cattle urine patches. *Agr. Forest Meteorol.* 156, 1-17.
- Leuning, R., Freney, J.R., Denmead, O.T., Simpson, J.R., 1985. A sampler for measuring atmospheric ammonia flux. *Atmos. Environ.* 19, 1117-1124.
- Lockyer, D.R., 1984. A system for the measurement in the field of losses of ammonia through volatilisation. *J. Sci. Food Agric.* 35, 37-848.
- Mosier, A., Kroeze, C., Nevison, C., Oenema, O., Seitzinger, S., van Cleemput, O., 1998. Closing the global  $\text{N}_2\text{O}$  budget: nitrous oxide emissions through the agricultural nitrogen cycle. *Nutr. Cycl. Agroecosyst.* 52, 225-248.
- Ni, K., Köster, J.R., Seidel, A., Pacholski, A., 2015. Field measurement of ammonia emissions after nitrogen fertilization—A comparison between micrometeorological and chamber methods. *Eur. J. Agron.* 71, 115-122.
- Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, B., Sirotenko, O., 2007. Agriculture. In *Climate Change 2007: Mitigation.*, in: Metz, B., Davidson, O.R., Bosch, P.R., Dave, R., Meyer, L.A. (Eds.), Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. .
- Suter, H., Sultana, H., Turner, D., Davies, R., Walker, C., Chen, D., 2013. Influence of urea fertiliser formulation, urease inhibitor and season on ammonia loss from ryegrass. *Nutr. Cycl. Agroecosyst.* 95, 175-185.
- Sutton, M.A., Milford, C., Dragosits, U., Place, C.J., Singles, R.J., Smith, R.I., Pitcairn, C.E.R., Fowler, D., Hill, J., ApSimon, H.M., Ross, C., Hill, R., Jarvis, S.C., Pain, B.F., Phillips, V.C., Harrison, R., Moss, D., Webb, J., Espenhahn, S.E., Lee, D.S., Hornung, M., Ullyett, J., Bull, K.R., Emmett, B.A., Lowe, J., Wyers, G.P., 1998. Dispersion, deposition and impacts of atmospheric ammonia: quantifying local budgets and spatial variability. *Environ. Pollut.* 102, 349-361.
- Zaman, M., Nguyen, M.L., Blennerhassett, J.D., Quin, B.F., 2008. Reducing  $\text{NH}_3$ ,  $\text{N}_2\text{O}$  and  $\text{NO}_3^-$ -N losses from a pasture soil with urease or nitrification inhibitors and elemental S-amended nitrogenous fertilizers. *Biol. Fertil. Soils* 44, 693-705.