

# Effective use of nitrogen fertilisers for growing garlic

Luji Bo<sup>1,2,3,4</sup>, Yingpeng Zhang<sup>1,2,3,4</sup>, Yan Li<sup>1,2,3,4,6</sup>, Jiafa Luo<sup>1,5</sup>, Ming Sun<sup>1,2,3</sup>, Ziwen Zhong<sup>1</sup>, Yongping Jing<sup>1,2,4</sup>

<sup>1</sup> Institute of Agricultural Resources and Environment, Shandong Academy of Agricultural Sciences, No.202 North Industrial Road, Jinan, China, 250100

<sup>2</sup> Key Laboratory of Agro-Environment of Huang-Huai-Hai Plain, Ministry of Agriculture, No.202 North Industrial Road, Jinan, China, 250100

<sup>3</sup> Shandong Provincial Key Laboratory of Agricultural Non-Point Source Pollution Control and Prevention, No.202 North Industrial Road, Jinan, China, 250100

<sup>4</sup> Shandong Provincial Engineering Research Center of Environmental Protection Fertilizers, No.202 North Industrial Road, Jinan, China, 250100

<sup>5</sup> Ruakura Research Centre, AgResearch Limited, Private Bag 3123, Hamilton, New Zealand, 3240

<sup>6</sup> Corresponding author: E-mail: [nkyliyan@126.com](mailto:nkyliyan@126.com)

## Abstract

An experiment was conducted on garlic (*Allium sativum* L.) to investigate the effects of nitrogen management on yield, economic benefit and the soil apparent nutrient balance, in the region of Laiwu town in Shandong province, China. The treatments included control (no N fertiliser), urea at 300 kg/ha, urea at 240 kg/ha, combined urea and commercial organic fertilisers at 120 kg N /ha each, and controlled-release nitrogen fertiliser at 192 kg N /ha. Results showed no significant difference both for garlic bulb yield and economic benefits between the urea treatments at different N application rates. The effect of the combined use of urea and organic fertiliser was similar to that of the urea application at the same N application rate. However, garlic bulb yields in the treatment of controlled-release nitrogen fertiliser were significantly higher than in the other fertiliser treatments, even the N application rate was lower. The net income from garlic in the treatment of controlled release fertiliser was also significantly higher than those in the other treatments ( $P < 0.05$ ). There was N surplus after the garlic growth season when urea was applied at 300 kg/ha, while there was N, P and K deficit when the controlled-release N fertiliser at 192 kg/ha was used to produce more garlic. It is suggested that use of controlled release N fertiliser combined with a supplement of other nutrients would be a sustainable strategy for fertiliser management in garlic production.

## Key words

Garlic bulb, Garlic bolt, N fertiliser application, Economic benefit, Nutrient balance

## Introduction

Garlic, classified under the *Alliaceae* family, is a widely consumed condiment vegetable. It has high nutritional value, and is rich in vitamins A and C. Garlic also contains antibiotic substances which makes it valuable for medical benefits (Bayan et al., 2014). Statistics in publications show the global importance of garlic, with about 19 million hectares planted and an overall annual production of nearly 327 million tonnes (Samavatean et al., 2011). About 80 thousand hectares of garlic are planted in China each year.

N fertilisers are often over-applied in an attempt to achieve high garlic yields. According to the survey, N fertiliser application rates in China are about 900-1500 kg/ha. A large amount of applied N has now accumulated in the soil due to this high level of N application, resulting in low N use efficiency. Furthermore, surplus N can be lost to the atmosphere and the groundwater (Savci, 2012). For example, the average concentration of nitrate in groundwater was up to 34.7 mg/L in the garlic production area in Jinxiang Town in China (Zhang et al., 2008).

The aims of this study were to examine the effects of nitrogen fertiliser management on garlic yield, economic benefit and the soil apparent nutrient balance. Our results will provide the direct evidence needed by farmers to apply N fertiliser for garlic production in an economically and environmentally rational manner.

## Materials and methods

### Experimental site

The field experiment was conducted on a typical vegetable production farm at Weiwangxu town, Laiwu city, Shandong Province, China (36°17' N, 117°27' E). The region has a typical warm and semi-humid continental monsoon climate. Mean annual temperature was 13.0 °C, and the lowest and highest mean monthly

temperatures were -14.5 °C in January and 36.7 °C in July, respectively. The average annual rainfall was 695 mm, concentrated in the summer. The study site was a field that had been under vegetable cultivation for decades. Soil samples from 0-20 cm depth showed the soil to be a Brown earth, with a pH (H<sub>2</sub>O) of 7.03, organic matter (OM) of 14.3 g/kg, and available N, P, and K of 156, 112, and 181 g/kg soil, respectively.

#### *Experimental design and crop management*

The field experiment consisted of five treatments, each in triplicate, with a randomized block design. Treatments included control (CK, no N fertiliser), urea at 300 kg/ha (FP), urea at 240 kg/ha (OPT), combined urea and commercial organic fertilisers at 120 kg/ha each (OM-CF), and controlled-release N fertiliser at 192 kg/ha (CRF) (Table 1). Fertilisers were applied three times in each of the FP, OPT and OM-CF treatments: 60% as a basal fertiliser, 25% as garlic bolt elongation fertiliser, and 15% as a garlic bulb expansion fertiliser (Table 1). In the Treatment CRF, thermoplastic resin-coated urea containing 4% coated material and 42% N, was applied only once as a basal fertiliser. Synthetic phosphate and potassium fertilisers were applied as basal fertilisers at rates of 90 kg P<sub>2</sub>O<sub>5</sub> /ha and 270 kg K<sub>2</sub>O /ha, respectively, in all the treatments except Treatment FP, which had a fertiliser application at a rate of 120 kg P<sub>2</sub>O<sub>5</sub> /ha and 255 kg K<sub>2</sub>O /ha. Garlic (*Allium sativum* L.) was cultivated in 7.0 m × 7 m field plots. Garlic seed was sown on 25<sup>th</sup> September 2012.

**Table 1. Fertiliser application in the experiment (kg/ha)**

Treatments	Base fertiliser			Supplemental fertiliser (Garlic bolt elongation)	Supplemental fertiliser (Garlic bulb expansion)	The total N applied
	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	N	N	
CK	0	90	270	0	0	0
FP	180	120	255	75	45	300
OPT	144	90	270	60	36	240
OM-CF	144	90	270	60	36	240
CRF	192	90	270	0	0	192

CK= no N fertiliser, FP= urea at 300 kg/ha, OPT= urea at 240 kg/ha, OM-CF= combined urea and commercial organic fertilisers at 120 kg/ha each, CRF= controlled-release nitrogen fertiliser at 192 kg/ha.

#### *Sampling and data analysis*


Garlic bolt and garlic bulb were harvested on 5<sup>th</sup> May 2013 and 28<sup>th</sup> May 2013, respectively. Methods of soil and agrochemical analysis were used for plant and soil analyses (Lu, 1999). Nutrient inputs included N, P, K from organic and inorganic fertilisers, and nutrient outputs included N, P, K uptaken by crops. The soil apparent nutrient balance was calculated by: soil apparent nutrient balance (kg/ha) = nutrient inputs (kg/ha) - nutrient outputs (kg/ha).

Statistical analyses were performed using SPSS 13.0 for Windows. An analysis of variance (ANOVA) test was used to determine the treatment effects on the measured variables. A least significant difference (LSD) multiple-comparison test was used to identify statistical differences among the treatments.

## **Results**

### *Effect of N application on the yield of garlic and garlic bolt*

The mean garlic bulb yields of each treatment are shown in Figure 1-A. Compared with no N application (CK), garlic bulb yields in the Treatment FP, OPT, CRF, and OM-CF increased by 31%, 29%, 55%, and 18%, respectively. All these increases were statistically significant ( $P < 0.05$ ). In addition, garlic bulb yields in Treatment CRF were 18%, 20%, and 30% higher than in the Treatments FP, OPT, and OM-CF, respectively. Compared with the yield from the Treatment OM-CF, the increases due to the application of CRF were significant ( $P < 0.05$ ), but there was no significant difference for garlic bulb yields among Treatment FP, OPT, and OM-CF. There was no significant difference for garlic bolt yields between Treatment CK and Treatment FP, OPT (Figure 1-B). Garlic bolt yields were lowest in Treatment CRF, while highest in Treatment OM-CF. The garlic bolt yields in Treatment OM-CF were 15%, 22%, 16%, and 35% higher than those in Treatment CK, FP, OPT, and CRF.



**Figure 1. Garlic bulb and garlic bolt yield in different fertilisation treatments. Different letters indicate significant differences ( $P < 0.05$ ).**

#### *Effect of N application on garlic economic benefits*

The cost of garlic seed, pesticide, and agricultural film was the same in all treatments. The lowest labour cost was in Treatment CRF (Table 2), as the fertiliser was only applied once as a basal fertiliser. The fertiliser cost in all treatments was in the following decreasing order: OM-CF > FP > OPT > CRF > CK. The total revenue and net income of garlic in all treatments followed the order of CRF > FP ≈ OPT ≈ OM-CF > CK, when the price of garlic in 2013 was considered. The net income of garlic in Treatment CRF was 61%, 20%, 22% and 35% higher than those in Treatment CK, FP, OPT, and OM-CF, respectively. The increase in income from the use of CRF was statistically higher than from the other fertilisers ( $P < 0.05$ ). The ratio of net income to cost was lowest in Treatment CK while highest in Treatment CRF. There was no significant difference in the ratio of net income to cost between Treatment CK and OM-CF, and between Treatment FP and OPT.

**Table 2. Average annual evaluation of economic benefit of garlic in different fertiliser treatments (costs and garlic price based on those in Shandong in 2013)**

Treatments	cost (Yuan/ha)			Total income (Yuan/ha)	Net income (Yuan/ha)	Net income/cost
	Seed, pesticide, agricultural film	Labor cost	Fertiliser			
CK	24000	15900	2424	128384 c	94160 c	3.75 c
FP	24000	15900	3369	162296 b	127127 b	4.61 b
OPT	24000	15900	3099	160350 b	125451 b	4.59 b
OM-CF	24000	15900	5149	150563 b	113613 b	4.07 c
CRF	24000	15150	3047	186915 a	152819 a	5.48 a

Values marked with different letters in the same column are significantly different ( $P < 0.05$ ).

#### *Effect of N application on soil apparent nutrient balance*

The N and P uptake in garlic was highest in Treatment CRF and lowest in Treatment CK (Table 3). There was no significant difference for nutrient uptake between Treatment FP and OPT. The apparent N deficiency in soil was 86 kg N/ha in Treatment CRF, while the apparent surplus N in soil was 50 and 21 kg N/ha in Treatment FP and OM-CF, respectively. Apparent P deficiency in soil occurred in all treatments. Apparent K deficit in soil was 20 kg K/ha in Treatment CRF, while the apparent surplus K from fertilisers in soil was 59, 25 and 10 kg/ha in Treatment CK, OPT and OM-CF, respectively.

**Table 3. Soil nutrient balance in different fertilization treatments**

Treatments	Nutrient input (kg/ha)			Nutrient output (kg/ha)			Nutrient net balance (kg/ha)		
	N	P	K	N	P	K	N	P	K
CK	0	39	224	191d	80d	165c	-191	-41	59
FP	300	52	212	250b	106ab	213b	50	-54	-1
OPT	240	39	224	240b	98bc	199b	0	-59	25
OM-CF	240	39	224	219c	89cd	214b	21	-50	10
CRF	192	39	224	278a	112a	244a	-86	-73	-20

CK= no N fertiliser, FP= urea at 300 kg/ha, OPT= urea at 240 kg/ha, OM-CF= combined urea and commercial organic fertilisers at 120 kg/ha each, CRF= controlled-release nitrogen fertiliser at 192 kg/ha

## Discussion

Compared with the nutrient level standard developed from the national soil survey, available nutrients in the soil in this study area were generally rich (National Soil Survey Office, 2002). The controlled-release N fertiliser had the best effect on garlic growth among all treatments, even when the N application rate was lowest. This reflects the high N use efficiency of controlled release N fertiliser, with the N released from the controlled-release fertiliser matching the N needed by garlic growth during critical periods (Zebarth et al., 2012). However, the garlic bolt yield in Treatment CRF experienced the lowest effect from the fertiliser, possibly due to insufficient N supply through the bolting period (a later period of growth). Therefore, the addition of N during that critical growth period could increase the yield of garlic bolt. The economic benefits in Treatment CRF was best, due to the low rate of N fertiliser application and labour costs.

Compared with conventional N fertilisation (ie. the Treatment FP), a 20% reduction of N use will not significantly affect garlic bulb and garlic bolt yield and its economic benefits. However, the ratio of net income to the cost of Treatment OM-CF was significantly lower than Treatment FP and OPT, due to the expensive of organic fertiliser. However, long-term organic fertiliser applications can maintain soil fertility and promote garlic yield (Zhang et al., 2013).

Nutrient deficiency and surpluses in soil depended on the input of fertilisers and nutrient uptake by garlic, which determined the soil sustainable production capacity. The results in the study indicated that conventional N application (ie. the treatment FP) could maintain garlic growth, resulting in a N surplus in soil. P deficit in soil was observed in all treatments, indicating that P fertiliser needs to be adequately increased to maintain an optimum P level in soil. It is noteworthy that the controlled-release N fertiliser can produce high yields of crop, at the expense of plant excessive uptake of soil nutrients. It is suggested that P and K fertilisers be applied when controlled release fertiliser is used to maintain soil nutrient balanced. Improvement of the controlled release fertiliser should supply N at the critical period of garlic growth.

## Conclusion

N surplus occurred in the conventional N application treatment when the N was applied at 300 kg N /ha. Compared with the conventional N application, a 20% reduction of N fertiliser did not significantly affect garlic yield and its economic benefits. A 36% reduction of N fertiliser, using controlled release N fertiliser, could increase garlic yield and its economic benefits, but would result in nutrient deficit soil which would affect long-term soil nutrient balance.

## Acknowledgements

This work was supported by the National Key Technology R&D Program of the Ministry of Science and Technology (2012BAD15B02), The Youth Scientific Research Foundation of Shandong Academy of Agricultural Sciences (2016YQN40), Key projects of scientific and technological innovation of Shandong Academy of Agricultural Sciences (2014CXZ01), and Taishan Scholars Program for Overseas Experts.

## References

- Bayan L, Koulivand PH and Gorji A (2014). Garlic: a review of potential therapeutic effects. *Avicenna Journal of Phytomedicine*. 4, 1-14.
- Lu R (1999). *Methods of Soil and Agrochemical Analysis*. China Agricultural Science and Technology Press, Beijing.
- National Soil Survey Office (2002). *China Soil*. China Agricultural Science and Technology Press, Beijing.
- Samavatean N, Rafiee S, Mobli H and Mohammadi A (2011). An analysis of energy use and relation between energy inputs and yield, costs and income of garlic production in Iran. *Renewable Energy*. 36, 1808-1813.
- Savci S (2012). Investigation of effect of chemical fertilizers on environment. *APCBEE Procedia*. 1, 287-292.
- Zebarth BJ, Snowdon E, Burton DL, Goyer C and Dowbenko R (2012). Controlled release fertilizer product effects on potato crop response and nitrous oxide emissions under rain-fed production on a medium-textured soil. *Canadian Journal of Soil Science*. 92, 759-769.
- Zhang W, Jiang Y, Meng P and Song S (2013). Effects of bio-organic fertilizers on development and yield of garlic. *Journal of Guizhou Agricultural Sciences*. 41, 63-65.
- Zhang W, Liu Z, Jiang L, Zheng F, Zhong Z, Wang M, Lin H and Song X (2008). Effect of nitrogen fertilization and optimal application N rates on garlic nutrition. *Chinese Agricultural Science Bulletin*. 24,

254-259.