

Wheat straw biochar reduces N₂O emission by increasing denitrification in alkaline and acidic submerged paddy soils

Jun Shan¹, Xu Zhao¹, Shutan Ma¹, Xiaoyuan Yan¹

¹ State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese Academy of Sciences, Nanjing 210008, China, Email: shanjun@issas.ac.cn or yanxy@issas.ac.cn

Abstract

Paddy fields are one of the most important N sinks in terrestrial ecosystems, and considerable N loss is caused by denitrification. Biochar has been recognized as useful soil amendment to paddy field in mitigating nitrous oxide (N₂O) emission. However, the key mechanisms responsible for the reduced N₂O emissions by biochar in paddy soils are still obscure. Here, using two paddy soils with contrasting pH, the denitrification and N₂O emission were investigated in soil amended with different amounts of biochar (0%, 0.5% and 5%) via soil slurry incubation combined with N₂/Ar technique. The results showed that biochar amendment significantly increased the pH values both in the alkaline and acidic soils. Biochar at 5% amendment rate significantly increased denitrification and significantly decreased N₂O emission in soils. In the alkaline soil, biochar at 0.5% amendment rate significantly increased denitrification, but had no effect on N₂O emission. Conversely, in the acidic soil, biochar at 0.5% amendment rate did not affect denitrification, but significantly reduced N₂O emission. The N₂O/(N₂+N₂O) ratio was significantly reduced by biochar amendment irrespective amendment rate both in alkaline and acidic soils. In the alkaline soil, biochar at 5% amendment rate significantly increased the abundance of *nosZ* genes, whereas biochar had no effect on the abundance of *nosZ* genes in the acidic soil irrespective of amendment rate. Our results suggested biochar effects in the alkaline soil were attributed to increase of denitrifying community, whereas biochar effects in the acidic soil was attributed to increase of pH.

Key Words

rice paddy, black carbon, denitrification, N₂/Ar technique, nitrous oxide

Introduction

Rice yield in China has increased substantially during the last three decades because of the rapidly increasing use of synthetic nitrogen (N) fertilizers (Ju et al., 2009). The N use efficiency in paddy field is generally low ranging from 30% to 40% (Zhu and Chen, 2002). The inefficiency of N use in the rice paddy field is mainly caused by denitrification, ammonia volatilization, and N leaching, among which denitrification is the predominant pathway for N loss way (Figure 1) (Xing and Zhu, 2000; Zhu and Chen, 2002). Also, substantial amounts of the applied synthetic N fertilizers in paddy soils are lost as nitrous oxide (N₂O) (Cai et al., 1997), a powerful greenhouse gas, which also contribute to the stratospheric ozone destruction (Ravishankara et al., 2009).




Figure 1. Sequential reductive pathway of denitrification with the location of enzymes i.e., nitrate reductase Nar (encoding *narG* and *napA* genes), nitrite reductase Nir (encoding *nirK* and *nirS* genes), nitric oxide reductase Nor (encoding *norB*, *norZ*, and *qNor* genes), and nitrous oxide reductase Nos (encoding *nosZ* gene).

It has been demonstrated that biochar application to paddy soils can significantly reduce emissions of N₂O in most previous studies (Zhang et al., 2010; Wang et al., 2011). Because in submerged paddy soils, N₂O is mainly produced through denitrification as an obligate intermediate product (Majumdar, 2013), the lower N₂O in the paddy soil amended with biochar can be attributed to a reduction in the total N denitrified or a stimulation of the reduction of N₂O to N₂ (Baggs, 2011; Cayuela et al., 2013). However, the key mechanisms responsible for the reduced N₂O emissions by biochar in paddy soils are still obscure. Also little is known about effects of biochar application on denitrification rate and N₂O/(N₂+N₂O) ratio in paddy soils.

In the present study, effects of biochar application on denitrification rate, N₂O emission and denitrifiers in one acidic and one alkaline paddy soils were investigated. We hypothesized that biochar application

increases denitrification rates both in acidic and alkaline paddy soils, consequently leading to reduced N₂O emissions. The abundance of denitrifier gene (*nosZ* gene) was also investigated to prove our hypothesis.

Methods

Soil and biochar

The alkaline and acidic rice paddy soils were collected from the plough layers (0–20 cm in depth) of fields under a rice–wheat rotation in Jiangdu (119° 42' E, 32° 35' N), Jiangsu province and Yingtian (28° 15' N, 116° 55' E), Jiangxi province, respectively. Biochar was produced from wheat straw in a muffle furnace under oxygen-limited conditions. Briefly, the starting pyrolysis temperature was set at 100 °C, subsequently elevated to 500 °C at 5 °C min⁻¹ and was held constant for 8 h.

Incubation experiment

All the incubation experiments were performed in 500 mL glass beaker, in which 150 g of soil (dry weight) were amended with different quantities of biochar resulting in three treatments: 0% biochar amendment (control), 0.5% biochar amendment, and 5% biochar amendment. The water content of all soil samples was adjusted to 60% water holding capacity (WHC). All the beakers were covered by parafilm with pin holes and incubated at 25 °C for 5 days, after which, soil pH, inorganic N contents of the soil-biochar matrix were determined. After 5 days aerobic incubation, subsamples (2.5 g) of the soil or soil-biochar matrix in the beakers was transferred into 12 mL glass tubes (Labco, UK) and 3 mL He-purged water was added, homogenized with the soil and stood for 5 min. Then another 6 mL He-purged water to result in soil slurry. For each treatment, forty glass tubes were prepared. Subsequently, all the tubes were sealed and preincubated underwater at 25 °C for 2 h. In the following 8 h incubations, eight replicate tubes at each sampling time (2 h interval) were removed and preserved with 200 µL saturated HgCl₂ solution, followed by centrifugation for 10 min (1800 g/25 °C). Then four replicate tubes were used to analyse dissolved N₂ by MIMS and N₂/Ar technique (Kana et al., 1994), while another four replicates were used to analyse dissolved N₂O by injecting 5 mL supernatant into 15 mL vacuum serum vials according to the method of Terry et al. (1981) and the N₂O concentration in the supernatant was determined by gas chromatography (GC). In our experiments, for simplicity, the denitrification rate was referred as the net N₂ flux because other processes which also generated N₂ play a minor role compared with denitrification. DNA was extracted from soil samples at the last time point for analysing functional gene abundances (*nosZ* gene) relevant to N₂O reduction by qPCR (Shan et al., 2016).

Data analysis

All statistical analyses were performed with software Sigmaplot 11.0 and the significant level was set at $P < 0.05$. The one way analysis of variance (ANOVA) followed by Fisher's least significant difference (LSD) test was performed to evaluate the differences among means of different biochar amendment treatments.

Results

Figure 2 presents the effects of biochar amendment (0%, 0.5%, and 5%) on pH and abundance of *nosZ* genes in the alkaline and acidic paddy soils. In the alkaline soil, 5% biochar amendment significantly increased soil pH values and size of *nosZ* gene abundance. Whereas, in the acidic soil, biochar amendment regardless of amendment rate significantly increased soil pH values and had no effect on size of *nosZ* gene abundance (Figure 2).




Figure 2. Effects of biochar amendment (0%, 0.5%, and 5%) on pH (A) and abundance of *nosZ* gene (B) in the alkaline and acidic paddy soils. Values are means with standard deviations ($n = 4$). The different letters above the bars indicate significant differences among the treatments.

Figure 3 shows the net N_2 flux, N_2O emission and $N_2O/(N_2+N_2O)$ ratio in the soils as affected by biochar amendment. In the alkaline soil, the net N_2 flux ($24.1 \text{ nmol } N_2\text{-N } g^{-1} h^{-1}$) was significantly increased by biochar amendment and the increase was 155.0% and 185.7% for 0.5% and 5% biochar amendment, respectively. The N_2O emission in the alkaline soil ($0.29 \text{ nmol } N_2O\text{-N } g^{-1} h^{-1}$) was significantly reduced by 5% biochar amendment ($0.15 \text{ nmol } N_2O\text{-N } g^{-1} h^{-1}$) and the $N_2O/(N_2+N_2O)$ ratio (1.2%) was significantly reduced by biochar amendment (0.47% and 0.33% for 0.5% and 5% biochar amendment, respectively) regardless of amendment rate (Figure 3). In the acidic soil, the net N_2 flux ($17.7 \text{ nmol } N_2\text{-N } g^{-1} h^{-1}$) was significantly increased by 5% biochar amendment ($25.4 \text{ nmol } N_2\text{-N } g^{-1} h^{-1}$) and the increase was 143.5%. The N_2O emission ($0.040 \text{ nmol } N_2O\text{-N } g^{-1} h^{-1}$) was significantly reduced by biochar amendment (0.0084 and $0.010 \text{ nmol } N_2O\text{-N } g^{-1} h^{-1}$ for 0.5% and 5% biochar amendment, respectively) irrespective of amendment rate. Similar to the alkaline soil, the $N_2O/(N_2+N_2O)$ ratio in the acidic soil (0.22%) was significantly reduced by biochar amendment (0.045% and 0.039% for 0.5% and 5% biochar amendment, respectively) regardless of amendment rate (Fig. 3).




Figure 3. Effects of biochar amendment (0%, 0.5%, and 5%) on denitrification, N₂O emission and ratio of N₂O to N₂ and N₂O in the alkaline and acidic paddy soils. Values are means with standard deviations ($n = 4$). The different letters above the bars indicate significant differences among the treatments.

Conclusion

Our results indicated that effects of biochar on denitrification and N₂O emission were soil-specific and depends on the biochar amendment rate. In the alkaline soil, biochar reduced N₂O emission by increasing denitrification being attributed to increase of *nosZ* gene transcription. Whereas, in the acidic soil biochar reduced N₂O emission by increasing denitrification being attributed to increase of pH.

References

- Baggs EM (2011): Soil microbial sources of nitrous oxide: recent advances in knowledge, emerging challenges and future direction. *Curr. Opin. in Env. Sust.* 3, 321-327.
- Cai ZC, Xing GX, Yan XY, Xu H, Tsuruta H, Yagi K, Minami K (1997): Methane and nitrous oxide emissions from rice paddy fields as affected by nitrogen fertilisers and water management. *Plant Soil* 196, 7-14.
- Cayuela ML, Sánchez-Monedero MA, Roig A, Hanley K, Enders A, Lehmann J (2013): Biochar and denitrification in soils: when, how much and why does biochar reduce N₂O emissions? *Sci. Rep.* 3, 1732
- Ju XT, Xing GX, Chen XP, Zhang SL, Zhang LJ, Liu XJ, Cui ZL, Yin B, Christie P, Zhu ZL, Zhang FS (2009): Reducing environmental risk by improving N management in intensive Chinese agricultural systems. *Proc. Natl. Acad. Sci. U. S. A.* 106, 3041-3046.
- Kana TM, Darkangelo C, Hunt, MD, Oldham JB, Bennett GE, Cornwell JC (1994): Membrane inlet mass spectrometer for rapid high-precision determination of N₂, O₂, and Ar in environmental water samples. *Anal. Chem.* 66, 4166-4170.
- Majumdar D (2013): Biogeochemistry of N₂O uptake and consumption in submerged soils and rice fields and implications in climate change. *Crit. Rev. Environ. Sci. Technol.* 43, 2653-2684.

- Ravishankara A, Daniel JS, Portmann RW (2009): Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century. *Science* 326, 123-125.
- Shan J, Zhao X, Sheng R, Xia Y, Ti C, Quan X, Wang S, Wei W, Yan, X (2016): Dissimilatory nitrate reduction processes in typical Chinese paddy soils: rates, relative contributions and influencing factors. *Environ. Sci. Technol.* (doi: 10.1021/acs.est.6b01765).
- Terry RE, Tate RL, Duxbury JM (1981): The effect of flooding on nitrous oxide emissions from an organic soil. *Soil Sci* 132:228–232.
- Wang JY, Zhang M, Xiong ZQ, Liu PL, Pan GX (2011): Effects of biochar addition on N₂O and CO₂ emissions from two paddy soils. *Biol. Fertil. Soils* 47, 887-896.
- Xing GX, Zhu ZL (2000): An assessment of N loss from agricultural fields to the environment in China. *Nutr. Cycl. Agroecosyst.* 57, 67-73.
- Zhang AF, Cui LQ, Pan GX, Li LQ, Hussain Q, Zhang XH, Zheng JW, Crowley D (2010): Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China. *Agric. Ecosyst. Environ.* 139, 469-475.
- Zhu Z, Chen D (2002): Nitrogen fertilizer use in China—contributions to food production, impacts on the environment and best management strategies. *Nutr. Cycl. Agroecosyst.* 63, 117-127.