




Understanding the variability in performance of the nitrification inhibitor 3,4-Dimethylpyrazole phosphate in Australian agricultural soils.

Helen Suter, Deli Chen (The University of Melbourne)
Charlie Walker (Incitec Pivot Fertilisers)




Background : Why Focus on Nitrification inhibitors?

➤ Reduce N₂O emissions



Australia's National GHG inventory (2014) (total CO₂-e)



93% =
agricultural
soils
= 18% of
agricultural
emissions


Fertiliser inputs
significant
contributor to N₂O

➤ Reduce nitrate leaching

➤ Improve nitrogen use efficiency

Background : Why Focus on 3,4-Dimethylpyrazole phosphate?

- Commercially available in Australia
- Potential for use across broad range of climates



3,4-Dimethylpyrazole phosphate

BUT

- Inconsistent results observed
- Reasons for this are unclear

How to address this? Experimental Methodology

➤ Laboratory Incubation experiment

➤ 30 soils, < 2 mm



➤ Treatments

- Control (no N)
- Fertiliser ($\text{NH}_4^+\text{-N}$)
- Fertiliser + DMPP

➤ $100 \mu\text{g NH}_4^+\text{-N / g soil} + 50 \mu\text{g NO}_3^-\text{-N / g soil}$



➤ 25°C , 60% WFPS, 28 days,

➤ Mineral N (2 M KCl 1:5) and N_2O collected



Results : Nitrification rate


Example of mineral N dynamic




Horsham soil (cropping) : 23% clay, 34% silt, 44% sand, pHw 8.5, OC 0.82%, N 0.08%

Results : Nitrification rate

Average nitrification rate (14 days) ($\mu\text{g NO}_3^- \text{N}$ produced/g soil/day)		
Treatment	Range	Average \pm standard error
Control	-4.61-26.89	1.37 \pm 0.99 ^a
Fertiliser	1.33-25.45	6.47 \pm 0.93 ^b
DMPP	-0.43-22.02	4.00 \pm 0.78 ^{ab}
Average net-nitrification rate (14 days) ($\mu\text{g NO}_3^- \text{N}$ produced/g soil/day)		
Fertiliser	-1.44-12.63	4.82 \pm 0.55 ^b
DMPP	-4.86-8.02	2.39 \pm 0.45 ^a



- Addition of fertiliser increases nitrification rate
- DMPP nitrification rate = control (no fertiliser)
- **Average 38% reduction in fertiliser induced nitrification rate with DMPP (range 9-100%)**





Other soil properties

- Clay (-ive) ($P < 0.05$)
- Mn (-ive) ($P < 0.1$)
- DTPA Cu (+ive)

fVnit_max: fraction of nitrification achieved with DMPP relative to the fertiliser only treatment

Results : N₂O emissions


Example of N₂O emission



Horsham soil (cropping) : 23% clay, 34% silt, 44% sand, pHw 8.5, OC 0.82%, N 0.08%


Results : N₂O emissions

Cumulative N ₂ O emissions (28 days) (µg N ₂ O-N/g soil)			
Treatment	range	Average ± standard error	Log ₁₀
Control	0.01-1.1	0.06±0.30	-1.20±0.64 ^a
Fertiliser	0.07-7.74	0.62±0.28	-0.59±0.45 ^b
DMPP	0.01-6.96	0.49±0.26	-0.92±0.64 ^{ab}
Cumulative Net-N ₂ O emissions (28 days) (µg N ₂ O-N/g soil)			
Fertiliser	-0.10-9.54	0.71±0.39	-0.57±0.48 ^a
DMPP	-0.01-10.75	0.52±0.35	-1.17±0.78 ^b



- Addition of fertiliser increases N₂O emissions
- DMPP nitrification rate = control (no fertiliser)

- **Average 55% reduction in fertiliser induced N₂O emissions with DMPP (range 0-100%)**



Other soil properties

- Mn (+ive) ($P < 0.05$)
- DTPA Zn (+ive) ($P < 0.05$)
- DTPA Fe (-ive) ($P < 0.01$)

- Effective tool for reducing nitrification and N₂O emissions across Australian agricultural soils
- High range of responses
- OM significantly ($P < 0.01$) affected the DMPP inhibition of nitrification
- pH significantly ($P < 0.01$) affected the DMPP inhibition of N₂O emissions
- Further investigation of the importance of properties other than organic matter and pH, and the role of soil trace elements and metals for their interactions with the inhibitor.
- The significant of the soil microbial community requires investigation
 - e.g. Bacterial versus archaeal responses

Thankyou

helencs@unimelb.edu.au

